

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

November 14, 2017

Matthew Brooks Regulatory Agent Director, Ag-Chem Consulting, LLC Agent for Ecostadt Technologies LLC 12644 Chapel Road, Suite 108 Clifton, VA 20124

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 – Update

Environmental Hazards Statements

Product Name: Ecoworks EC

EPA Registration Number: 89152-4 Application Date: October 31, 2017 OPP Decision Number: 535393

Dear Dr. Brooks:

The U.S. Environmental Protection Agency (EPA) is in receipt of your application for notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Biopesticides and Pollution Prevention Division (BPPD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records. You must submit one (1) copy of the final printed labeling with the modifications.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Page 2 of 2 EPA Reg. No. 89152-4 OPP Decision No. 535393

If you have any questions, please contact James Parker of my team by phone at (703) 306-0469 or via email at parker.james@epa.gov.

Sincerely,

Anchew L. kupcelon

Andrew Bryceland, Team Leader Biochemical Pesticides Branch Biopesticides and Pollution Prevention Division (7511P) Office of Pesticide Programs

Enclosure

89152-4

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

11/14/2017

[Note: Texts in [ ] are optional language]

ECOWORKS EC October 31, 2017 – Version 1

## ECOWORKS EC (EPA Reg. # 89152- 4 ) MASTER LABEL

Sub Label A: Agriculture crops, Greenhouses, Turf &

Ornamental Use

Sub Label B: Residential/Home Owner & Residential Greenhouse

Use

Sub Label C: Residential/ Home Owner Use- Single Dose vial

Sub Label D: Residential/Home Owner Use- Single Dose sachet

### **ECOWORKS EC**

[Additional Brand Names- ECOHALT EC, NEEM WORKS EC, ECOMAXIMA EC]
BOTANICAL INSECTICIDE, MITICIDE, NEMATICIDE & FUNGICIDE

FOR USE ON VEGETABLES, FRUITS, NUTS, COMMERCIAL ROW CROPS, CEREALS, COMMERCIAL GREENHOUSES, SHADE HOUSES, INTERIORSCAPES, INDOOR & OUTDOOR ORNAMENTALS, NURSERIES, TURF, ORNAMENTALS, TREES SHRUBS, HOME OWNER USES AND RESIDENTIAL GREENHOUSES.

FOR THE CONTROL OF INSECTS, MITES, NEMATODES AND FUNGAL DISEASES.
PRODUCT CAN BE USED UP TO DAY OF HARVEST IN ALL FOOD AND NON-FOOD CROPS.

 ACTIVE INGREDIENT: Cold Pressed Neem oil [\*]
 70.00%

 OTHER INGREDIENTS:
 30.00%

 100.00%

[\* Contains 2200 PPM Azadirachtin]

## KEEP OUT OF REACH OF CHILDREN CAUTION

#### **FIRST AID**

#### If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- · Call a poison control center or doctor for treatment advice.

#### If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

#### If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

#### If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call your poison control center at 1-800-222-1222. For information on product, usage, call 1-888-858-7378, Seven days a week, 6:30 am to 4.30 pm Pacific Time. During other times, call the poison control center 1-800-222-1222.

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:

ECOSTADT TECHNOLOGIES LLC 1024 Iron Point Road, Suite 100-1318

Folsom, CA 95630 Phone: 916-357-6607 Net Contents: 2, 4, 8, 16, 32 OZ, 1, 2.5, 5 Gallons

EPA Reg. # 89152-4

EPA Est. # Batch No.:

[Note: Texts in [ ] are optional language]

## ECOWORKS EC (EPA Reg. # 89152- 4 ) MASTER LABEL

Sub Label A: Agriculture crops, Greenhouses, Turf & Ornamental Use

### **ECOWORKS EC**

[Additional Brand Names- ECOHALT EC, NEEM WORKS EC, ECOMAXIMA EC] BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE & FUNGICIDE

FOR USE ON VEGETABLES, FRUITS, NUTS, COMMERCIAL ROW CROPS, CEREALS, COMMERCIAL GREENHOUSES, SHADE HOUSES, INTERIORSCAPES, INDOOR & OUTDOOR ORNAMENTALS, NURSERIES, TURF, ORNAMENTALS, TREES AND SHRUBS, FOR THE CONTROL OF INSECTS, MITES, NEMATODES AND FUNGAL DISEASES. PRODUCT CAN BE USED UP TO DAY OF HARVEST IN ALL FOOD AND NON-FOOD CROPS.





[\*Contains 2200 PPM Azadirachtin]

# KEEP OUT OF REACH OF CHILDREN CAUTION

|                         | FIRST AID   |  |  |  |  |  |  |
|-------------------------|---|--|--|--|--|--|--|
| If on skin or clothing: | <ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>   |  |  |  |  |  |  |
| If inhaled:             | <ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>                                     |  |  |  |  |  |  |
| If in eyes:             | <ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |  |  |  |  |  |  |
| If swallowed:           | <ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul> |  |  |  |  |  |  |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call your poison control center at 1-800-222-1222. For information on product, usage, etc., call 1-888-858-7378, Seven days a week, 6:30 am to 4.30 pm Pacific Time. During other times, call the poison control center 1-800-222-1222.

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:

ECOSTADT TECHNOLOGIES LLC

1024 iron Point Road, Suite 100-1318

Folsom, CA 95630

Phone: 916-357-6607

Net Contents: 2, 4, 8, 16, 32 OZ, 1, 2.5, 5

Gallons

EPA Reg. # 89152-4

EPA Est. No.: Batch No:

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before use.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

Long-sleeved shirt and long pants:

Chemical-resistant (such as barrier laminate, butyl, nitrile, neoprene, polyvinyl chloride, or viton) gloves.

Socks and shoes

Follow manufacturer's instructions for cleaning /maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **USER SAFETY RECOMMENDATIONS**

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### **ENVIRONMENTAL HAZARDS**

For Terrestrial Uses Only. This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage, ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any irrigation system unless the chemigation instructions on this label are followed. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

Long-sleeved shirt and long pants
Chemical – resistant (such as barrier laminate, butyl, nitrile, neoprene, polyvinyl chloride, or viton) gloves
Socks and shoes

#### NON-AGRICULTURAL USE REQUIREMENTS

These requirements apply to uses of this product that are NOT within the WPS for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For other uses including golf courses and other non-agricultural uses, do not enter treated areas without protective clothing until sprays have dried.

#### PRODUCT INFORMATION

ECOWORKS EC is an emulsifiable concentrate containing cold pressed Neem oil 70%. ECOWORKS EC controls insects such as worms, thrips, aphids, white flies, leaf miners, beetles, mites and nematodes etc. ECOWORKS EC repels and/or disrupts feeding and growth of the target insect pests.

ECOWORKS EC is effective for the prevention and control of the following fungal diseases: powdery mildew, downy mildew, rust, leaf spot, scab, blight, anthracnose, dollar spot, and molds.

#### **COMPATIBILITY**

ECOWORKS EC has been found to be compatible with most commonly used insecticides, fungicides and fertilizers. However, to avoid plant damage, a jar compatibility test must be conducted before mixing (or) applying this product. To conduct a jar compatibility test, take a pint of water and adjust pH between 5.5 to 7.0 using a buffering agent (Buffering agents include products such as Buffer PS, Buffer Xtra Strength etc.). To test tank compatibility of ECOWORKS EC and pesticide formulations including water soluble, emulsifiable, microemulsion, suspension concentrate and surfactants, first mix ECOWORKS EC @1

TSP/Pint in 5.5 to 7.0 pH adjusted water, mix well, and then add intended pesticides at labelled rates and mix well again.

#### **TANK MIXING**

For tank mix compatibility tests of ECOWORKS EC and pesticides formulations including dry wettable powders and dry water dispersible granules, first mix labelled rates of dry products in a Pint of 5.5 to 7.0 pH adjusted water, mix well, and then add ECOWORKS EC @ 1 TSP, and mix well again. In case tank mixing involves 3 categories of products including ECOWORKS EC, liquid formulations (such as water soluble, emulsifiable, microemulsion, suspension concentrates & surfactants) and dry products (wettable powder and water dispersible granules), first mix dry products in pH 5.5-6.0 adjusted Pint water followed by ECOWORKS EC and liquid products. After thorough mixing of ECOWORKS EC and intended tank mixing pesticides, let the solution stand for 15-30 minutes. If separation, sedimentation, flocculation and grainy appearance in the spray mixture is formed, it is not compatible and such mixtures may not be used. The tank mixture is considered compatible in case there is no separation, sedimentation, flocculation and grainy appearance in the mixture and such tank mixtures may be used. Once tank mixture compatibility is established, test spray mixtures on a small sample of plants for possible adverse effects for phytotoxic effects.

As a general recommendation, do not use ECOWORKS EC with products such as Bordeaux mixture, Triphenyltin hydroxide, lime sulfur, Rayplex Iron (or) other high alkaline insecticides, fungicides and fertilizers. If ECOWORKS EC is mixed with very mild alkaline products (pH 7.5-8.0) that show tank mixing compatibility, use such tank mixtures within 8 hours.

Always follow directions for use, precautions and limitations for use on all product labels used in combination. Always follow manufacturer's directions for Use and Precautionary Statements.

#### **APPLICATION INSTRUCTIONS**

#### READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USE

ECOWORKS EC is exempt from tolerances and may be applied as directed to any food or non-food crop up to and including the day of harvest at a rate not exceeding 14 Pints (20 grams active ingredient) per acre per application.

#### **GENERAL PRECAUTIONS**

Do not apply ECOWORKS EC on just germinated, newly transplanted, wilted and drought stressed plants and plants with very tender leaves.

Do not apply ECOWORKS EC during hot temperature conditions (>90 F). Apply this product during early morning and late evening time to prevent potential leaf burns and plant injuries.

Exercise extreme care when combining ECOWORKS EC with oil and surfactant based products.

Do not apply to sensitive plant species such as poinsettias, some rose species, impatiens, hibiscus flowers, carnations, and comice pears.

Waxy bloom on certain ornamental plants may be reduced after an application.

Applications may remove 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

MIXING AND STORAGE: Shake well before mixing. Use spray water with ambient conditions (20-35 °C or 68-95 °F) for mixing and avoid using cold water. In case full spray water is not available at suggested ambient conditions, premix product with a few gallons of water at 20-35 °C or 68-95 °F and then mix with cold water. Use spray mixtures immediately (or) within 24 hours of preparation. The recommended pH range is between 5.5 and 7.0 for ECOWORKS EC alone (or) when used with other products. When mixing with other approved agrichemicals, always ensure proper agitation in the spray tank to ensure uniform application. If required, use a buffering agent.

Refer to the Use Rate tables below to determine the amount of ECOWORKS EC for a required area to be treated. First, add half the volume of water your proposing to dilute the product in based on the Use Rate table to the mix tank, fill half water in mix tank, add determined amount of ECOWORKS EC and then add remaining water for proper mixing.

ECOWORKS EC disperses freely when added to water. For proper dilution and uniform distribution on plant canopy, ensure proper agitation in mixing tanks (or) vessels. When mixing with other agrochemicals, add solid constituents (such as wettable powders, water dispersible granules or micronutrients) last in the form of a slurry.

**APPLICATION METHOD AND EQUIPMENT:** ECOWORKS EC can be applied as a foliar spray (or) soil drench (or) soil-less media (e.g., greenhouses and mushroom houses) to control listed insects, mites, nematodes and fungal diseases. Soil drenches can be used to control listed soil-borne pests, including soil-borne larvae of foliar insect pests. When applying as a drench, avoid excessive leaching. ECOWORKS EC can be applied through sub-surface soil treatment equipment (e.g. turf grass). To repel adult flies, apply through fogging equipment.

Apply ECOWORKS EC using any powered (or) manual pesticide application equipment, including high-volume, low- volume, ultra-low volume, electrostatic, fogging and aerial applications. Follow original manufacturer's recommendations when using these types of equipment.

For optimum results, make 2 to 3 applications at 7 to 10 day intervals. Ensure thorough foliar coverage for effective pest and disease control.

#### **ECOWORKS EC USE SITES:**

Use ECOWORKS EC on the following crops:

#### FIELD AND GREENHOUSE FOOD CROPS-

Brassica (cole) crops, cucurbits, eggplants, herbs and spices, legumes, peppers, tomatoes, and other miscellaneous crops grown in greenhouses.

**Edible Fungi Group-**Agaricus, cremini, enoki, maitake, oyster, portobello, shitake, and white mushrooms.

**Root and Tuber Vegetables Group-** Artichokes, beets, carrots, ginger, horseradish, potatoes, radishes, rutabagas, sweet potatoes, turmeric, turnips, yams.

Leafy and Brassica Crop Vegetables Crop Grouping (including Brassica Leafy Vegetables) - Amaranth, broccoli, Brussels sprouts, cabbage, cauliflower, celery, chervil,

Chinese cabbage, collards, cress, endives, fennel, kale, kohlrabi, lettuce, mizuna, mustard greens, parsley, purslane, rape greens, rhubarb, spinach, Swiss chard.

**Legume Vegetables Group -** Beans (field, kidney etc.), chickpeas, cowpeas, guar, jack beans, lablab beans, lentils, peas, pigeon peas, soybeans, sword beans.

**Fruiting Vegetables Group -** Eggplants, ground cherries, pepinos, peppers, pimentos, tomatillos, tomatoes.

**Cucurbit Vegetables Group -** Bitter melons, Chayotes, Chinese wax gourds, citron melons, cucumbers, gherkins, gourds, muskmelons (such as cantaloupes, casabas Crenshaw etc.), pumpkins, squash, watermelons.

**Citrus Fruits Group -** Calamondins, citrus citrons, citrus hybrids, grapefruits, kumquats, lemons, limes, mandarins, oranges, pummelos, Satsuma mandarins.

Pome Fruits Group - Apples, crab apples, loquats, mayhaws, oriental pears, pears, quinces.

Stone Fruits Group - Apricots, cherries, nectarines, peaches, plums, prunes

**Berries Group -** Blackberries and caneberries, blueberries, currants, elderberries, gooseberries, huckleberries, loganberries, raspberries, strawberries, youngberries.

Forage - Alfalfa, clover, trevoil or vetch.

**Cereal Grains Group -** Barley, buckwheat, corn, millet, oats, popcorn, rice, rye, sorghum, teosintes, triticale hybrids, wheat, wild rice.

#### **Cotton and Tobacco**

Herbs and Spices Group - Allspice, angelica, anise, annatto, balm, basil, black and white peppers, borage, burnet, chamomile, caper buds, cardamom, caraway, cassia, catnip, celery seeds, chervil, chives, cinnamon, clary, cloves, coriander (cilantro), costmary, cumin, curry leaf, dills, fennels, fenugreek, grains of paradise, horehound, hyssop, juniper berry, lavender, lemongrass, lavage, mace, marigolds, marjoram, mustard seeds, nasturtium, nutmeg, parsley, pennyroyal, poppy seeds, rosemary, rue, saffron, sage, savory, sweet bay (bay leaf), tansy, tarragon, thyme, vanilla, wintergreen, woodruff, wormwood.

Bulb Vegetables (Allium spp.) Crop Group- Garlic, leeks, onions, shallots

**Nuts Group -** Almonds, beechnuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapin, filberts, hickory nuts, lychee nuts, macadamias, pecans, pistachios, walnuts.

**Oilseed Group -** Canola, castor, crambe, guar, jojoba, peanuts, rape, safflower, sesame, soybean, sunflower.

**Tropical fruits -** Atemoyas, bananas, breadfruits, cherimoyas, durians, guavas, malangas, mangos, papaya, passion fruits, star fruits.

**Miscellaneous Food and Non-Food Crops -** Asparagus, avocados, birdseed, cacao, coffee, edible flowers, feijoa, figs, ginseng, grapes, guayule, hops, kiwis, okras, olives, palms, papayas,

pawpaws, persimmons, pineapples, rambutans, sugarcane, tamarillos, tea, water chestnuts, watercress.

**ORNAMENTAL PLANTS -** African violets, ageratum, aster, aucuba, begonia, cacti, calendula, calla, carnation, ceanothus, chrysanthemum, cineraria, coleus, cotoneaster, cyclamen, daffodil, dahlia, delphinium, ficus, foliage plants, fuchsia, gardenia, geranium, gloxinia, hyacinth, hydrangea, iris, ivy, lily, maidenhair fern, marigold, narcissus, orchid, pansy, pelargonium, peony, phlox, pittosporum, poinsettia, pyracantha, rubber plant, snapdragon, stock, tulip, wandering jew, yew, yucca, zinnia.

#### CHRISTMAS TREES AND CHRISTMAS TREE PLANTATIONS

**ORNAMENTAL TREES and SHRUBS -** Andromeda, arbor vitae, ash, Austrian pine, azalea, beech, birch, birds nest spruce, blue spruce, bougainvillea, boxwood, butternut, camellia, cedar, chamaecyparis, cherry, crabapple, Cyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honeylocust, horse chestnut, ilex, juniper, larch laurel, lilac, linden, London plane, magnolia, manvilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach pine, photinia, planetree, pines, poplar, privet, quince, rhododendron, roses, spruce, sycamore, white cedar, and white pine.

**TURF and TURFGRASS** – Bent grass, Bermuda grass, bluegrass, centipede grass, fescue, ryegrass, St. Augustine, wheatgrass, zoysia grass.

**INTERIORSCAPES, SOIL-LESS MEDIUM CROPS - (Non-Residential)** Vegetables, Herbs, Ornamentals.

**NON-CROP USES-** Manures, manure piles, composts, compost piles, poultry litter, poultry litter piles, cull piles, biochar, mulches, potting soil, granular fertilizers, top soil, crop waste dumps, farm yards, fuel storage areas, fence rows, rights of way, fallow land and barrier strips.

# (1). ECOWORKS EC USES FOR VEGETABLES, FRUITS, NUTS, COMMERCIAL ROW CROPS, CEREALS OUTDOOR ORNAMENTALS, TURF, ORNAMENTALS, TREES AND SHRUBS- FOLIAR APPLICATIONS

ECOWORKS EC is intended for use for outdoor plants and food crops, mushrooms grown in polyhouses, plants grown indoors or in greenhouses, interiorscapes and nurseries. It can be used to control any of the following insects: mites, nematodes and listed fungal diseases.

Use tables below to determine appropriate use rate for your site / pest combination. Rates are provided in pints of ECOWORKS EC. When combining with other insecticides, use half recommended rate of ECOWORKS EC listed below. Use a spray water volume of 25 to 100 gallons depending on crop and crop stages. Do not exceed 14 pints/acre.

| ECOWORKS USE RATES FOR OUTDOOR PLANTS INCLUDING: FOOD CROPS AND |  |  |  |  |  |  |  |
|---|--|--|--|--|--|--|--|
| NON-CROP USE (TREES, TU   | NON-CROP USE (TREES, TURFGRASS, NURSERY & ALL OUTDOOR ORNAMENTAL |  |  |  |  |  |  |
|   | PLANTS)  |  |  |  |  |  |  |
| PESTS PINTS /ACRE* REMARKS                                      |  |  |  |  |  |  |  |

| WHITEFLIES Greenhouse whiteflies, Silver leaf whiteflies, Woolly whiteflies  | Low use -1 Pint<br>Mid use - 2 Pints<br>High use- 4 Pints | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
|--|---|---|
| LEAF MINERS Azalea, leaf miners, Birch leaf miners, Citrus leaf miners, Serpentine leaf miners, Vegetable leaf miners. | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| SCALES Brown soft scales, California red scales, Coffee scales, Olive scales, San Jose scales                          | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| MEALY BUGS Citrus mealybugs  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| THRIPS Citrus thrips, Onion thrips, Thrips palmi,  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| APHIDS Cotton aphids, Green peach aphids, Pea aphids, Potato aphids  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| PSYLLIDS<br>Pear psylla  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| LEAFHOPPERS Grape leafhoppers  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| BUGS Boxelder bugs, Chinch bugs, Lygus bugs Spittle bugs, Stink bugs   | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear.  Thorough coverage is recommended. |
| FLIES Blueberry maggots, Cherry maggots, Crane flies, Fruit flies, Midges, Onion maggots, Walnut husk flies            | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| SAWFLIES European pine sawflies, Yellow headed pine sawflies   | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints     | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |

| CATERPILLARS  Armyworms, Artichoke plume moths, Bag worms, Boll worms, Bud worms, Cabbage butterflies, Cabbage Loopers, Canker worms, Case worms, Corn ear worms, Cut worms, Diamond-backed moths, Fruit worms, Grape leaf skeletonizers, Gypsy moths, Hickory shuck worms, Hornworms, Imported cabbage worms Leaf perforators, Leaf rollers, Melon worms, Navel Orange worms, Oblique banded leaf rollers, Omnivorous leaf rollers. Oriental fruit moths, Pickle worms, Pine tip moths, Pinworms, red-banded leaf rollers, Sod webworms, Soybean loopers, Tent caterpillars, Tobacco budworms, Tussock moths. | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Repeat applications 7-14 day intervals Apply early when pests appear. Thorough coverage is recommended.   |
|--|---|---|
| BEETLES Bark Beetles, Blueberry flea beetles, Boll weevils, Colorado potato beetles, Flea beetles, Japanese beetles, Leaf beetles, Mexican bean beetles, Pepper weevils, Phylloxera Rose Chafers, Twig girdlers.   | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| WEEVILS Black vine weevils, Strawberry vine weevils  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| BORERS Peach twig borers, Peachtree borers, Dogwood borers, Cranberry borers.  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| MOLE CRICKETS  | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.  |
| NEMATODES Burrowing nematodes, Dagger nematodes, Golden nematodes, Root knot nematodes   | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Apply in sufficient amount of water to penetrate in the soil to a depth of 12 inches. Repeat applications every 3 or 4 weeks or as needed.  Thorough coverage is recommended. |

| MITES Red spider mites, Brown mite, Clover mites, Conifer spider mite, European red mites, Spruce spider mites and Two spotted spider mites | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.    |
|---|---|---|
| Diseases powdery mildew, downy mildew, rust, black spot, leaf spot, botrytis, anthracnose, blight, dollar spot, mold etc.                   | Low use 1 Pint<br>Mid use 2 Pints<br>High use 4 Pints | Repeat applications 7-14 day intervals. Apply early when diseases appear. Thorough coverage is recommended. |

<sup>\*</sup>When pest and disease infestation is heavy or when plant canopy is dense, use mid to high recommended rates on the table. When combining with other insecticides & fungicides, use 1-2 pints/acre with label rates of other products.

## (2). ECOWORKS EC USES FOR COMMERCIAL GREENHOUSES, SOIL-LESS MEDIUM, SHADECLOTH AND NURSERIES- FOLIAR APPLICATIONS

Use the table below to determine appropriate use rate for each pest. Foliar sprays for individual plants should thoroughly wet both sides of the leaves without causing runoff.

## ECOWORKS EC USE RATES FOR PLANTS GROWN INDOORS OR IN GREENHOUSES, SOIL-LESS MEDIUM CROPS, UNDER SHADECLOTHES & NURSERIES

| PESTS                  | PINTS/<br>100 GALLONS* | REMARKS                                       |
|------------------------|------------------------|---|
| WHITEFLIES             | 2 -4 Pints             | Repeat applications 7-14 day intervals. Apply |
| Greenhouse whiteflies, | for 50, 000 sq. ft     | early when pests appear.                      |
| Silver leaf whiteflies |                        | Thorough coverage is recommended.             |
| LEAF MINERS            | 2 -4 Pints             | Repeat applications 7-14 day intervals. Apply |
| Serpentine leaf miners | for 50,000 sq. ft.     | early when pests appear.                      |
|                        |                        | Thorough coverage is recommended.             |
| SOFT SCALES            | 2 -4 Pints             | Repeat applications 7-14 day intervals. Apply |
|                        | for 50,000 sq. ft.     | early when pests appear.                      |
|                        |                        | Thorough coverage is recommended.             |
| MEALY BUGS             | 2 -4 Pints             | Repeat applications 7-14 day intervals. Apply |
|                        | for 40,000 sq. ft.     | early when pests appear.                      |
|                        |                        | Thorough coverage is recommended.             |
| THRIPS                 | 2-4 Pints              | Repeat applications 7-14 day intervals. Apply |
| Western flower thrips  | for 40,000 sq. ft.     | early when pests appear.                      |
|                        |                        | Thorough coverage is recommended.             |
| APHIDS                 | 2 -4 Pints             | Repeat applications 7-14 day intervals. Apply |
| Green peach aphids,    | for 40,000 sq. ft.     | early when pests appear.                      |
| Pea aphids, Cotton     |                        | Thorough coverage is recommended.             |
| aphids, Rose aphids    |                        |   |
| LACEWINGS              | 2 -4 Pints             | Repeat applications 7-14 day intervals. Apply |

| Azalea lacewings  | for 40,000 sq. ft.               | early when pests appear. Thorough coverage is recommended.   |
|---|----------------------------------|--|
| FLIES Crane flies, Fungus gnats, Shore flies  | 2 -4 Pints<br>for 40,000 sq. ft. | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.     |
| CATERPILLARS Army worms, Bag worms, Cut worms, Leaf hoppers, Leaf rollers, Loopers, Spruce budworms, Webworms                               | 2 -4 Pints<br>for 40,000 sq. ft. | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.     |
| BORERS Peachtree borers   | 2 -4 Pints<br>for 50,000 sq. ft. | Repeat applications 7-14 day intervals.  Apply early when pests appear.  Thorough coverage is recommended.   |
| BEETLES Bark beetles, Flea beetles, Japanese beetles  | 2 -4 Pints<br>for 50,000 sq. ft. | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.     |
| WEEVILS Black vine weevils, strawberry vine weevils   | 2 -4 Pints<br>for 40,000 sq. ft. | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.     |
| NEMATODES Burrowing nematodes, Dagger nematodes, Golden nematodes, Root knot nematodes  | 2-4 Pints<br>for 40,000 sq. ft.  | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.     |
| MITES Red spider mites, Brown mite, Clover mites, Conifer spider mite, European red mites, Spruce spider mites and Two spotted spider mites | 2-4 Pints<br>for 40,000 sq. ft.  | Repeat applications 7-14 day intervals. Apply early when pests appear. Thorough coverage is recommended.     |
| Diseases Powdery mildew, downy mildew, rust, black spot, leaf spot, botrytis, anthracnose, alternaria etc.                                  | 2-4 Pints<br>for 40,000 sq. ft.  | Repeat applications 7-14 days intervals. Apply early when diseases appear. Thorough coverage is recommended. |

<sup>\*</sup> Use 2 Pints under low to medium pest pressure and 4 Pints when high infestation is observed. When combining with other insecticides & fungicides, use 1-2 pints/acre with label rates of other products. Do not exceed 14 pints/acre.

## (3). ECOWORKS EC USES FOR COMMERCIAL GREENHOUSES, SOIL-LESS MEDIUM, SHADECLOTH AND NURSERIES- FOLIAR LOW VOLUME APPLICATIONS

# ECOWORKS EC SPRAY PREPARATION FOR LOW VOLUME APPLICATIONS FOR ANY PLANTS GROWN INDOORS, GREENHOUSE, UNDER SHADECLOTHES, SOIL-LESS MEDIUM CROPS AND NURSERIES

| PESTS       | RATE/<br>1 GAL   | RATE/<br>5 GAL           | RATE/<br>10 GAL           | RATE/<br>15 GAL         | RATE/<br>20 GAL           | REMARKS  |
|-------------|------------------|--------------------------|---------------------------|-------------------------|---------------------------|--|
| WHITEFLIES  | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4<br>Fl. Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |
| LEAF MINERS | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |
| SOFT SCALES | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |
| MEALY BUGS  | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |
| THRIPS      | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |
| APHIDS      | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |
| LACE WINGS  | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |
| FLIES       | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days. |

| CATERPILLARS       | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days.    |
|--------------------|------------------|--------------------------|---------------------------|-------------------------|---------------------------|---|
| BORERS             | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days.    |
| BEETLES            | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days.    |
| WEEVILS            | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days.    |
| NEMATODES          | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days.    |
| MITES              | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when pests first appear. Repeat every 5 to 7 days.    |
| FUNGAL<br>DISEASES | 10 ml<br>(2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 100 ml<br>(3.4 Fl.<br>Oz) | 150 ml<br>(5 Fl.<br>Oz) | 200 ml<br>(6.8 Fl.<br>Oz) | Spray when diseases first appear. Repeat every 5 to 7 days. |

When infestation is heavy or when plant canopy is dense, use rates up to twice (2X) shown in table, but do not exceed 14 pints/ acre. When combining with other insecticides & fungicides, use half rates shown in table.

## (4). ECOWORKS EC USES FOR GREENHOUSES, SOIL-LESS MEDIUM, SHADECLOTH AND NURSERIES-RECIRCULATORY APPLICATIONS

Use ECOWORKS EC in re-circulatory applications for effective control of foliar and soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or thrips and diseases, etc.

Measure required quantity of product as per table provided below and add to a bucket of water and mix it well prior to application to re-circulatory system applications.

## ECOWORKS EC APPLICATION TABLE FOR ALL INDOOR PLANTS SUCH AS GREENHOUSE, UNDER SHADECLOTHES, INTERIORSCAPES, NURSERIES, SOIL-LESS MEDIUM CROPS AND RECIRCULATORY APPLICATIONS

| Gallon of  | А                        | APPLICATION                |                         |                            |             |
|------------|--------------------------|----------------------------|-------------------------|----------------------------|-------------|
| water      | 0.05%                    | 0.1%                       | 0.15%                   | 0.2%                       | INTERVAL    |
| 1 GALLON   | 2 ml<br>(2/5 Tsp)        | 4 ml<br>(4/5 Tsp)          | 6 ml<br>(1.2 Tsp)       | 8 ml<br>(1.6 Tsp)          | 7 – 14 days |
| 5 GALLONS  | 10 ml<br>(2 Tsp)         | 20 ml<br>(4 Tsp)           | 30 ml<br>(1 Fl.<br>Oz)  | 40 ml<br>(1.4 Fl.<br>Oz)   | 7 – 14 days |
| 10 GALLONS | 20 ml<br>(4 Tsp)         | 40 ml<br>(1.4 Fl.<br>Oz)   | 60 ml<br>(2 Fl.<br>Oz)  | 80 ml<br>(2.7 Fl.<br>Oz)   | 7 – 14 days |
| 20 GALLONS | 40 ml<br>(1.4 Fl.<br>Oz) | 80.0 ml<br>(2.7 Fl.<br>Oz) | 120 ml<br>(4 Fl.<br>Oz) | 160 ml<br>(5.4 Fl.<br>Oz)  | 7 – 14 days |
| 30 GALLONS | 60 ml<br>(2 Fl. Oz)      | 120 ml<br>(4 Fl. Oz)       | 180 ml<br>(6 Fl. Oz)    | 240 ml<br>(8 Fl. Oz)       | 7 – 14 days |
| 40 GALLONS | 80 ml<br>(2.7 Fl. Oz)    | 160 ml<br>(5.4 Fl. Oz)     | 240 ml<br>(8 Fl. Oz)    | 320 ml<br>(10.8 Fl.<br>Oz) | 7 – 14 days |

Use ECOWORKS EC in re-circulatory system for control of listed foliar pests, soil borne insect larvae, fungus gnats, nematodes& fungal diseases. Only dilute ECOWORKS EC at 0.05 to 0.1 %. Do not use higher concentration without checking for plant phytotoxicity and mixing compatibility. Agitate re-circulation mixture until product is completely dispersed.

## (5). ECOWORKS EC USES FOR GREENHOUSES, SOIL-LESS MEDIUM, SHADECLOTH AND NURSERIES- DRENCH APPLICATIONS

Use ECOWORKS EC as a soil drench for effective control of foliar and soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or thrips, diseases, etc.

Measure required quantity of product as per table below, add to a bucket of water and mix it well prior to drench applications.

ECOWORKS EC APPLICATION TABLE FOR DRENCH APPLICATIONS FOR INDOOR PLANTS AND OUTDOOR PLANTS IN GREENHOUSES, SOIL-LESS MEDIUM CROPS

| Gallon of | Al           | APPLICATION  |            |                  |             |
|-----------|--------------|--------------|------------|------------------|-------------|
| water     | 0.05%        | 0.01%        | 0.15%      | 0.2%             | INTERVAL    |
| 1 GALLON  | 2 ml         | 4 ml         | 6 ml       | 8 ml             | 7 – 14 days |
|           | (2/5 Tsp)    | (4/5 Tsp)    | (1.2 Tsp)  | (1.6 Tsp)        |             |
| 5 GALLON  | 10 ml        | 20 ml        | 30 ml      | 40 ml            | 7 – 14 days |
|           | (2 Tsp)      | (4 Tsp)      | (1 Fl.     | (1.4 Fl.         |             |
|           |              |              | Oz)        | Oz)              |             |
| 10 GALLON | 20 ml        | 40 ml        | 60 ml      | 80 ml            | 7 – 14 days |
|           | (4 Tsp)      | (1.4 Fl.     | (2 Fl.     | (2.7 Fl.         |             |
|           |              | Oz)          | Oz)        | Oz)              |             |
| 20 GALLON | 40 ml        | 80.0 ml      | 120 ml     | 160 ml           | 7 – 14 days |
|           | (1.4 Fl.     | (2.7 Fl.     | (4 Fl.     | (5.4 Fl.         |             |
|           | Oz)          | Oz)          | Oz)        | Oz)              |             |
| 30 GALLON | 60 ml        | 120 ml       | 180 ml     | 240 ml           | 7 – 14 days |
|           | (2 Fl. Oz)   | (4 Fl. Oz)   | (6 Fl. Oz) | (8 Fl. Oz)       |             |
| 40 GALLON | 80 ml        | 160 ml       | 240 ml     | 320 ml           | 7 – 14 days |
|           | (2.7 Fl. Oz) | (5.4 Fl. Oz) | (8 Fl. Oz) | (10.8 Fl.<br>Oz) |             |

Drench soil in the pot with one (1) pint of finished product dilution per gallon of soil. Use lower concentration (0.05%) for mild infestations and higher concentration (0.2%) for severe infestation.

#### (5). ECOWORKS EC USES IN AERIAL APPLICATIONS

For aerial applications in crops, turf, ornamentals and trees, apply 4-8 pints/acre diluted in 10-20 gallons of water. Avoid application when conditions do not support uniform coverage or when excessive drift may occur. Do not apply directly to animals, animal houses, aquatic animals or aquatic systems.

Do not apply this product in any type of agricultural chemigation systems.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Do not store this product above 100 degrees F or below 20 degrees F for extended periods of time. Store in original container in a dry, cool, well-ventilated area.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed on site or at an approved waste disposal facility.

**Container Disposal:** Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container ¼ full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

#### **IMPORTANT: PLEASE READ BEFORE USE**

#### LIMITED WARRANTY

ECOSTADT TECHNOLOGIES LLC warrants that to the extent permitted by applicable law, (a) this product conforms to chemical description on its label; (b) this product is reasonably fit for purposes stated on its label, subject to inherent risks referred to herein, when used in accordance with its directions; and (c) that directions, cautions and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. To the extent permitted by applicable law, there are no express warranties other than those set forth herein. ECOSTADT TECHNOLOGIES LLC neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. To the extent permitted by applicable law, ECOSTADT TECHNOLOGIES LLC expressly excludes and disclaims all implied warranties of merchantability, fitness for purpose (or) any other warranty of quality of performance.

To the extent permitted by applicable law, this warranty does not extend to, and user shall be solely responsible for, any loss or damage that results from the use of this product in any manner that is inconsistent with this label's directions, or cautions.

To the extent permitted by applicable law, user's exclusive remedy and ECOSTADT TECHNOLOGIES LLC's or seller's exclusive liability for any claim loss, damage, or injury resulting from use or handling of this product, whether (or) not based in contract, negligence, strict liability in tort, (or) otherwise, shall be limited, at ECOSTADT TECHNOLOGIES LLC's option, to replacement, (or) repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

### [Optional marketing claims, may be used on any panel]

- Controls listed Insects, Mites and Nematodes.
- Insecticide, Miticide & Nematicide
- Broad Spectrum Control.
- Can be used the day of harvest.
- Rose and flower spray.
- Garden vegetable and Ornamental spray.
- Tomato and vegetable spray.
- Foliar and systemic application.

[Note: Texts in [ ] are optional language]

### ECOWORKS EC (EPA Reg. #. 89152-4)

Sub Label B: Residential/Home Owner & Residential Greenhouse Use

### **ECOWORKS EC**

[Additional Brand Names- ECOHALT EC, NEEM WORKS EC, ECOMAXIMA EC] BOTANICAL INSECTICIDE, MITICIDE, NEMATICIDE & FUNGICIDE

FOR HOME GARDEN AND RESIDENTIAL GREENHOUSE USE FOR THE CONTROL OF INSECTS, MITES, NEMATODES AND FUNGAL DISEASES PRODUCT CAN BE USED UP TO DAY OF HARVEST IN ALL FOOD AND NON-FOOD CROPS.





[\* Contains 2200 PPM Azadirachtin]

# KEEP OUT OF REACH OF CHILDREN CAUTION

#### **FIRST AID**

#### If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

#### If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

#### If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call your poison control center at 1-800-222-1222. For information on product, usage, call 1-888-858-7378, Seven days a week, 6:30 am to 4.30 pm Pacific Time. During other times, call the poison control center 1-800-222-1222.

Medical: Hazard information Services (H.I. S1) 800-228-5635, Extension 169.

ECOSTADT TECHNOLOGIES LLC

1024 Iron Point Road, Suite 100-1318 Net Contents: 2, 4, 8, 16, 32 OZ, 1 Gal, 2.5 Gal, 5 Gallons

Folsom, CA 95630 EPA Reg.# 89152-4

EPA Est. #

Phone: 916-357-6607 Batch No.:

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before use.

#### **ENVIRONMENTAL HAZARDS**

For Terrestrial Uses Only. This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage, ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### PRODUCT INFORMATION

ECOWORKS EC is an emulsifiable concentrate containing 70% cold pressed Neem oil. ECOWORKS EC controls worms, thrips, aphids, white flies, leaf miners, beetles, mites and nematodes. ECOWORKS EC repels and disrupts feeding and growth of the listed insect pests. ECOWORKS EC is effective for the prevention and control of the following fungal diseases: powdery mildew, downy mildew, rust, leaf spot, scab, blight, anthracnose, dollar spot, molds.

#### **APPLICATION INSTRUCTIONS**

#### READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USE

ECOWORKS EC is exempt from tolerances and may be applied as directed to any food or non-food crop up to and including day of harvest at a rate not exceeding 14 Pints (20 grams active ingredient) per acre per application.

#### **GENERAL PRECAUTIONS**

Do not apply ECOWORKS EC on just germinated, newly transplanted, wilted and drought stressed plants and plants with tender leaves.

Do not apply ECOWORKS EC during hot temperature conditions (>90 F). Apply this product during early morning and late evening time to prevent potential leaf burns and plant injuries.

Do not apply this product to to sensitive plant species such as poinsettias, some rose species, impatiens, hibiscus flowers, carnations, Comice pears etc.

Waxy bloom on certain ornamental plants may be reduced after an application.

Applications may remove 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

MIXING AND STORAGE: Shake well before mixing. Use spray water with ambient conditions (20-35 °C or 68-95°F) for mixing and avoid using cold water. In case full spray water is not available at suggested ambient conditions, premix product with a few gallons of water at 20-35 °C or 68-95°F and then mix with cold water. Use spray mixtures immediately (or) within 24 hours of mixing. The recommended pH range is between 5.5 and 6.5 for ECOWORKS EC alone (or) when used with other products. If required, use a buffering agent. When mixing with other approved agrichemicals, always ensure proper agitation in the spray tank to ensure uniform application.

#### (1). ECOWORKS EC USES IN HOME GARDEN PLANTS- FOLIAR APPLICATIONS

For foliar applications, apply ECOWORKS EC with hand operated or low volume power sprayer equipment. Follow equipment manufacturer's instructions/recommendations when making applications.

To Mix- Measure required product quantity (see table below) into a clean bucket the add required amount of water and then stir well before loading into a sprayer.

| ECOWORKS EC APPLICATION RATES FOR HOME GARDEN CROPS SUCH AS VEGETABLES, HERBS, SPICES, FRUITS, BERRIES AND ORNAMENTAL HOME GARDEN CROPS |  |              |       |                   |             |  |  |  |
|---|--|--------------|-------|-------------------|-------------|--|--|--|
| PESTS & DILUTION RATE APPLICATION   |  |              |       |                   |             |  |  |  |
| CROP  | CROP  DISEASES SUCH AS  Quart Water  Quart Gallon Water  One Gallon Water  INTERVALS |              |       |                   |             |  |  |  |
| Leafy Vegetables including but not  | Leafrollers,<br>Loopers,   | 2.5 ml<br>(½ | 10 ml | 50 ml<br>(1.7 Fl. | 7 – 14 days |  |  |  |

| limited to: Broccoli, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach  | Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, rust  | Tsp)                   | ( 2 Tsp)          | Oz)                      |             |
|---|---|------------------------|-------------------|--------------------------|-------------|
| Root Vegetables including but not limited to: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yam                               | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>rust | 2.5 ml<br>(½<br>Tsp)   | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Fruiting Vegetables including but not limited to: Eggplant, Pepper, Tomatillo, Tomato   | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites Powdery mildew, Downy mildew, rust                             | 2.5 ml<br>(1/2<br>Tsp) | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Cucurbit Vegetables including but not limited to: Cucumber, Gourd, Muskmelon, Pumpkin, Squash, Watermelon, Cantaloupe, Casaba, Gherkins, Melons, Zucchini | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites<br>Powdery<br>mildew,<br>Downy mildew,          | 2.5 ml<br>( ½<br>Tsp)  | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Legume Vegetables including but not limited to: Bean, Chickpea, Lentil, Pea   | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites   | 2.5 ml<br>( ½<br>Tsp)  | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |

|  | Powdery<br>mildew,<br>Downy mildew,<br>rust   |                       |                   |                          |             |
|--|---|-----------------------|-------------------|--------------------------|-------------|
| Bulb Vegetables including but not limited to: Garlic, Onion, Shallot   | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew, | 2.5 ml<br>(½<br>Tsp)  | 10 ml<br>(2 Tsp)  | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Berries including but<br>not limited to:<br>Blackberry,<br>Blueberry,<br>Raspberry,<br>Strawberry,<br>Gooseberry | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Anthracnose, Botrytis   | 2.5 ml<br>( ½<br>Tsp) | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Herbs and Spices including but not limited to: Chive, Dill, Fennel, Mustard, Sage, Basil, Coriander, Peppermint  | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew, | 2.5 ml<br>( ½<br>Tsp) | 10 ml<br>(2 Tsp)  | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Nut trees including<br>but not limited to:<br>Almond, Walnut,<br>Pecan, Pistachio                                | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,  | 2.5 ml<br>(½<br>Tsp)  | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Pome fruits including<br>but not limited to:<br>Apple, Pear (check<br>plant safety for<br>Comice Pear variety    | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,  | 2.5 ml<br>( ½<br>Tsp) | 10 ml<br>(2 Tsp)  | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |

| before application)  | Thrips, Mites,  |                       |                   |                          |             |
|--|---|-----------------------|-------------------|--------------------------|-------------|
| belore application)  | Powdery mildew, Downy mildew, Anthracnose, Botrytis   |                       |                   |                          |             |
| Stone fruits including<br>but not limited to:<br>Apricot, Cherry,<br>Nectarine, Peach,<br>Plum | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Anthracnose, Botrytis                               | 2.5 ml<br>( ½<br>Tsp) | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Citrus fruits including<br>but not limited to:<br>Grapefruit, Lemon,<br>Orange                 | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>Anthracnose,<br>Botrytis | 2.5 ml<br>(½<br>Tsp)  | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Ornamental Plants including but not limited to: Chrysanthemum, Coleus, Roses, Poinsettia       | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>Blackspot, rust          | 2.5 ml<br>( ½<br>Tsp) | 10 ml<br>(2 Tsp)  | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |
| Ornamental Trees & Shrubs  | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery   | 2.5 ml<br>( ½<br>Tsp) | 10 ml<br>( 2 Tsp) | 50 ml<br>(1.7 Fl.<br>Oz) | 7 – 14 days |

| mildew,         |  |  |
|-----------------|--|--|
| Downy mildew,   |  |  |
| Blackspot, rust |  |  |
|                 |  |  |

# (2). ECOWORKS EC USE FOR HOME GARDEN PLANTS, INDOOR AND OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, INTERIORSCAPES, AND SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS- RE-CIRCULATORY APPLICATIONS

Use ECOWORKS EC in re-circulatory applications for effective control of foliar and soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or thrips, and listed fungal diseases.

Measure required quantity of the product as provided in the table below into a bucket of water and mix it well prior to addition to re-circulatory system applications.

# OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, RESIDENTIONAL INTERIORSCAPES, AND RESIDENTIONAL SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS- RECIRCULATORY APPLICATIONS

| Gallon of  | А         | APPLICATION |           |           |             |
|------------|-----------|-------------|-----------|-----------|-------------|
| water      | 0.05%     | 0.1%        | 0.15%     | 0.2%      | INTERVAL    |
| 1 GALLON   | 2 ml      | 4 ml        | 6 ml      | 8 ml      | 7 – 14 days |
|            | (2/5 Tsp) | (4/5 Tsp)   | (1.2 Tsp) | (1.6 Tsp) |             |
| 5 GALLONS  | 10 ml     | 20 ml       | 30 ml     | 40 ml     | 7 – 14 days |
|            | (2 Tsp)   | (4 Tsp)     | (1 Fl.    | (1.4 Fl.  |             |
|            |           |             | Oz)       | Oz)       |             |
| 10 GALLONS | 20 ml     | 40 ml       | 60 ml     | 80 ml     | 7 – 14 days |
|            | (4 Tsp)   | (1.4 Fl.    | (2 Fl.    | (2.7 Fl.  |             |
|            |           | Oz)         | Oz)       | Oz)       |             |
| 20 GALLONS | 40 ml     | 80.0 ml     | 120 ml    | 160 ml    | 7 – 14 days |
|            | (1.4 Fl.  | (2.7 Fl.    | (4 Fl.    | (5.4 Fl.  |             |
|            | Oz)       | Oz)         | Oz)       | Oz)       |             |
| 30 GALLONS | 60 ml     | 120 ml      | 180 ml    | 240 ml    | 7 – 14 days |
|            | (2 Fl.    | (4 Fl.      | (6 Fl.    | (8 Fl.    |             |
|            | Oz)       | Oz)         | Oz)       | Oz)       |             |
| 40 GALLONS | 80 ml     | 160 ml      | 240 ml    | 320 ml    | 7 – 14 days |
|            | (2.7 Fl.  | (5.4 Fl.    | (8 Fl.    | (10.8 Fl. |             |

| Oz) | Oz) | Oz) | Oz) |  |
|-----|-----|-----|-----|--|
|     |     |     |     |  |

<sup>\*</sup> Use ECOWORKS EC in re-circulatory system for the control of listed foliar pests, soil borne insect larvae, fungus gnats, nematodes & fungal diseases etc. Only dilute ECOWORKS EC at 0.05 to 0.1%. Do not use higher concentration without checking for plant phytotoxicity and mixing compatibility. Agitate re-circulation mixture until product is completely dispersed. Do not exceed 14 Pints/Acre.

# (3). ECOWORKS USES FOR HOME GARDEN PLANTS, INDOOR AND OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, INTERIORSCAPES, AND SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS- DRENCH APPLICATIONS

Use ECOWORKS EC as a soil drench for effective control of foliar and soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or thrips, and listed fungal diseases.

Measure required quantity of the product as provided in the table below and add to a bucket of water and mix it well prior to application to re-circulatory system applications.

#### ECOWORKS EC APPLICATION TABLE FOR HOME & GARDENS DRENCH APPLICATIONS FOR INDOOR AND OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, INTERIORSCAPES, AND SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS

| Gallon of | Gallon of AMOUNT OF ECOWORKS EC* |           |            |                 |             |  |
|-----------|----------------------------------|-----------|------------|-----------------|-------------|--|
| water     | 0.05%                            | 0.01%     | 0.15%      | 0.2%            | INTERVAL    |  |
| 1 GALLON  | 2 ml                             | 4 ml      | 6 ml       | 8 ml            | 7 – 14 days |  |
|           | (2/5 Tsp)                        | (4/5 Tsp) | (1.2 Tsp)  | (1.6 Tsp)       |             |  |
| 5 GALLON  | 10 ml                            | 20 ml     | 30 ml      | 40 ml           | 7 – 14 days |  |
|           | (2 Tsp)                          | (4 Tsp)   | (1 Fl. Oz) | (1.4 Fl.<br>Oz) |             |  |
| 10 GALLON | 20 ml                            | 40 ml     | 60 ml      | 80 ml           | 7 – 14 days |  |
|           | (4 Tsp)                          | (1.4 Fl.  | (2 Fl.     | (2.7 Fl.        |             |  |
|           |                                  | Oz)       | Oz)        | Oz)             |             |  |
| 20 GALLON | 40 ml                            | 80.0 ml   | 120 ml     | 160 ml          | 7 – 14 days |  |
|           | (1.4 Fl.                         | (2.7 Fl.  | (4 Fl.     | (5.4 Fl.        |             |  |
|           | Oz)                              | Oz)       | Oz)        | Oz)             |             |  |
| 30 GALLON | 60 ml                            | 120 ml    | 180 ml     | 240 ml          | 7 – 14 days |  |
|           | (2 Fl.                           | (4 Fl.    | (6 Fl.     | (8 Fl.          |             |  |
|           | Oz)                              | Oz)       | Oz)        | Oz)             |             |  |
| 40 GALLON | 80 ml                            | 160 ml    | 240 ml     | 320 ml          | 7 – 14 days |  |
|           | (2.7 Fl.                         | (5.4 Fl.  | (8 Fl.     | (10.8 Fl.       |             |  |
|           | Oz)                              | Oz)       | Oz)        | Oz)             |             |  |

\*Drench soil in the pot with one (1) pint of finished product dilution per gallon of soil. Use lower concentration (0.05%) for mild infestations and higher concentration (0.2%) for severe infestation.

#### STORAGE AND DISPOSAL

#### Do Not Contaminate Water, Food, or Feed by Storage or Disposal.

Store in a cool dry place out of reach of children and pets and away from direct sunlight, windows, flames and other sources of heat or ignitions. Keep container tightly closed and in original container when not in use. Do not store product such that it is exposed to ultraviolet light (sunlight) or moisture.

Non-refillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

**If partly filled**: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### **IMPORTANT: PLEASE READ BEFORE USE**

#### LIMITED WARRANTY

ECOSTADT TECHNOLOGIES LLC warrants that to the extent permitted by applicable law (a) this product conforms to chemical description on its label; (b) this product is reasonably fit for purposes stated on its label, subject to inherent risks referred to herein, when used in accordance with its directions; and (c) that directions, cautions and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. To the extent permitted by applicable law, there are no express warranties other than those set forth herein. ECOSTADT TECHNOLOGIES LLC neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. To the extent permitted by applicable law, ECOSTADT TECHNOLOGIES LLC expressly excludes and disclaims all implied warranties of merchantability, fitness for purpose (or) any other warranty of quality of performance.

To the extent permitted by applicable law, this warranty does not extend to, and user shall be solely responsible for, any loss or damage that results from the use of this product in any manner that is inconsistent with this label's directions, or cautions.

To the extent permitted by applicable law, user's exclusive remedy and ECOSTADT TECHNOLOGIES LLC's or seller's exclusive liability for any claim loss, damage, or injury resulting from use or handling of this product, whether (or) not based in contract, negligence, strict liability in tort, (or) otherwise, shall be limited, at ECOSTADT TECHNOLOGIES LLC's option, to replacement, (or) repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

#### [Optional marketing claims, may be used on any panel]

- Controls listed Insects, Mites and Nematodes.
- Insecticide, Miticide & Nematicide
- Broad Spectrum Control.
- Can be used the day of harvest.
- Rose and flower spray.
- Garden vegetable and Ornamental spray.
- Tomato and vegetable spray.
- Foliar and systemic application.

[Note: Texts in [ ] are optional language]

## ECOWORKS EC (EPA Reg. #. 89152-4) Sub Label C: Residential/Home Owner Use

### **ECOWORKS EC**

[Additional Brand Names- ECOHALT EC, NEEM WORKS EC, ECOMAXIMA EC]
BOTANICAL INSECTICIDE, MITICIDE, NEMATICIDE & FUNGICIDE
READY TO MAKE PRODUCT (RTM)
PREMEASURED SINGLE DOSE DELIVERY VIAL

FOR HOME GARDEN AND RESIDENTIAL GREENHOUSE USE.
FOR THE CONTROL OF INSECTS, MITES, NEMATODES AND FUNGAL DISEASES.
PRODUCT CAN BE USED UP TO DAY OF HARVEST IN ALL FOOD AND NON-FOOD CROPS.





| ACTIVE INGREDIENT: Cold Pressed Neem oil [*] | 70.00%  |
|--|---------|
| OTHER INGREDIENTS:                           | 30.00%  |
|  | 100.00% |

[\* Contains 2200 PPM Azadirachtin]

# KEEP OUT OF REACH OF CHILDREN CAUTION

#### **FIRST AID**

#### If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

#### If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

#### If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.

#### Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call your poison control center at 1-800-222-1222. For information on product, usage, call 1-888-858-7378, Seven days a week, 6:30 am to 4.30 pm Pacific Time. During other times, call the poison control center 1-800-222-1222.

Medical: Hazard information Services (H.I. S1) 800-228-5635, Extension 169.

ECOSTADT TECHNOLOGIES LLC 1024 Iron Point Road, Suite 100-1318

Folsom, CA 95630 Phone: 916-357-6607 Net Contents: 5 vials (Total Volume: 50ml, 3 1/3 Tbs), 10 vials (Total Volume: 100ml, 6 2/3 Tbs), 20 vials (Total Volume: 200ml, 13 1/3 Tbs), 40 vials (Total Volume: 400ml, 26 2/3Tbs).

Each Vail contains 10 ml (2/3 Tbs)

EPA Reg. # 89152-4

EPA Est. # Batch No.:

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before use.

#### **ENVIRONMENTAL HAZARDS**

For Terrestrial Uses Only. This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage, ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### PRODUCT INFORMATION

ECOWORKS EC is an emulsifiable concentrate containing 70% cold pressed Neem oil. ECOWORKS EC controls insects such as worms, thrips, aphids, white flies, leaf miners, beetles, mites and nematodes etc. ECOWORKS EC repels, disrupts feeding and growth of insect pests.

ECOWORKS EC is effective for the prevention and control of fungal diseases such as powdery mildew, downy mildew, rust, leaf spot, scab, blight, anthracnose, dollar spot, molds etc.

#### **APPLICATION INSTRUCTIONS**

#### READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USE

ECOWORKS EC is exempt from tolerances and may be applied as directed to any food (or) non-food crop up to and including day of harvest at a rate not exceeding 14 Pints (20 grams active ingredient) per acre per application.

#### **GENERAL PRECAUTIONS**

Do not apply ECOWORKS EC on just germinated, newly transplanted, wilted and drought stressed plants and plants with tender leaves.

As followed in oil based formulations, do not apply ECOWORKS EC during hot temperature conditions (>90 F). It is recommended to apply during early morning and late evening time to prevent potential leaf burns and plant injuries.

Exercise care when combining with oil and surfactant based products.

It is recommended not to apply to sensitive plant species such as poinsettias, some rose species, impatiens, hibiscus flowers, carnations, Comice pears etc.

Waxy bloom on certain ornamental plants may be reduced after an application.

Applications may remove 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

**MIXING AND STORAGE:** Shake well before mixing. Always use this product promptly after mixing with water. Use spray water at ambient conditions (20-35 35 °C or 68-95°F) for mixing and avoid using cold waters. In case spray water is cold, first premix the product with a few gallons of water at 20-35 35 °C or 68-95°F and then mix with cold water. Use spray mixture immediately (or) within few hours of mixing.

#### (1). ECOWORKS EC USES FOR HOME GARDEN PLANTS- FOLIAR APPLICATIONS

For foliar applications, apply ECOWORKS EC with hand operated or power sprayer equipment for low volume. Follow equipment manufacturer recommendations when making applications.

Open ECOWORKS EC packet containing single dose vials. Drop required number of vials in to hand sprayer as per table below, fill water, shake well and spray.

## ECOWORKS EC FOLIAR APPLICATION RATES FOR HOME GARDEN PLANTS SUCH AS VEGETABLES, HERBS, SPICES, FRUITS, BERRIES AND ORNAMENTALS

|   | PESTS &   |                        | ECOWORI                 | APPLICATION            |   |
|---|---|------------------------|-------------------------|------------------------|---|
| CROP  | DISEASES<br>SUCH AS   | One<br>Gallon<br>water | Five<br>Gallon<br>water | Ten<br>Gallon<br>water | INTERVALS   |
| Leafy Vegetables including but not limited to: Broccoli, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery | 1 vial                 | 5 vials                 | 10 vials               | 7 - 14 days.  Spray early as soon as pest infection is noticed. |

|   | mildew, Downy mildew, rust  |        |         |          |   |
|---|---|--------|---------|----------|---|
| Root Vegetables including but not limited to: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yam                               | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>rust | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Fruiting Vegetables including but not limited to: Eggplant, Pepper, Tomatillo, Tomato   | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites<br>Powdery<br>mildew,<br>Downy mildew,<br>rust  | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Cucurbit Vegetables including but not limited to: Cucumber, Gourd, Muskmelon, Pumpkin, Squash, Watermelon, Cantaloupe, Casaba, Gherkins, Melons, Zucchini | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites<br>Powdery<br>mildew,<br>Downy mildew,          | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Legume Vegetables including but not limited to: Bean, Chickpea, Lentil, Pea   | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites<br>Powdery<br>mildew,<br>Downy mildew,<br>rust  | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Bulb Vegetables including but not   | Leafrollers,<br>Loopers,  | 1 vial | 5 vials | 10 vials | 7 - 14 days.<br>Spray early as                                  |

| limited to:<br>Garlic, Onion, Shallot  | Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,                             |        |         |          | soon as pest<br>infection is<br>noticed.                        |
|--|---|--------|---------|----------|---|
| Berries including but<br>not limited to:<br>Blackberry,<br>Blueberry,<br>Raspberry,<br>Strawberry,<br>Gooseberry                     | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Anthracnose, Botrytis   | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Herbs and Spices including but not limited to: Chive, Dill, Fennel, Mustard, Sage, Basil, Coriander, Peppermint                      | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew, | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Nut trees including<br>but not limited to:<br>Almond, Walnut,<br>Pecan, Pistachio  | t trees including Leafrollers, not limited to: Loopers, Armyworms,  |        | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Pome fruits including<br>but not limited to:<br>Apple, Pear (check<br>plant safety for<br>Comice Pear variety<br>before application) | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew,                         | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |

|  | Anthracnose,   |        |         |          |   |
|--|--|--------|---------|----------|---|
|  | Botrytis   |        |         |          |   |
| Stone fruits including<br>but not limited to:<br>Apricot, Cherry,<br>Nectarine, Peach,<br>Plum | ts including Leafrollers, Loopers, Armyworms,  |        | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Citrus fruits including<br>but not limited to:<br>Grapefruit, Lemon,<br>Orange                 | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Anthracnose, Botrytis                      | 1 vial | 5 vials | 10 vials | 7- 14 days.  Spray early as soon as pest infection is noticed.  |
| Ornamental Plants including but not limited to: Chrysanthemum, Coleus, Roses, Poinsettia       | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>Blackspot, rust | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Ornamental Trees & Shrubs  | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>Blackspot, rust | 1 vial | 5 vials | 10 vials | 7 - 14 days.  Spray early as soon as pest infection is noticed. |

## (2). ECOWORKS EC USES FOR HOME GARDEN PLANTS, INDOOR AND OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, INTERIORSCAPES, AND SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS- DRENCH APPLICATION

Use ECOWORKS EC as a soil drench for effective control of foliar and soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or thrips, fungal diseases, etc.

Open ECOWORKS EC packet containing single dose vials. Drop required number of vials into a clean bucket with water as prescribed in the table below and stir the bucket well.

#### ECOWORKS EC APPLICATION TABLE FOR HOME & GARDENS DRENCH APPLICATIONS FOR INDOOR AND OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, INTERIORSCAPES, AND SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS

| Gallons of | AM       | APPLICATION |          |          |             |
|------------|----------|-------------|----------|----------|-------------|
| water      | 0.05%    | 0.1%        | 0.15%    | 0.2%     | INTERVAL    |
| 5 GALLONS  | 1 vial   | 2 vials     | 3 vials  | 4 vials  | 7 – 14 days |
| 10 GALLONS | 2 vials  | 4 vials     | 6 vials  | 8 vials  | 7 – 14 days |
| 20 GALLONS | 4 vials  | 8 vials     | 12 vials | 16 vials | 7 – 14 days |
| 30 GALLONS | 6 vials  | 12 vials    | 18 vials | 24 vials | 7 – 14 days |
| 40 GALLONS | 8 vials  | 16 vials    | 24 vials | 32 vials | 7 – 14 days |
| 50 GALLONS | 10 vials | 20 vials    | 30 vials | 40 vials | 7 – 14 days |

<sup>\*</sup>Drench soil in the pot with one (1) pint of finished product dilution per gallon of soil. Use lower concentration (0.05%) for mild infestations and higher concentration (0.2%) for severe infestation.

#### STORAGE AND DISPOSAL

#### Do Not Contaminate Water, Food, or Feed by Storage or Disposal.

Store in a cool dry place out of reach of children and pets and away from direct sunlight, windows, flames and other sources of heat or ignitions. Keep containers tightly closed and in original containers when not in use. Do not store product such that it is exposed to ultraviolet light (sunlight) or moisture.

Non-refillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

**If partly filled**: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### **IMPORTANT: PLEASE READ BEFORE USE**

#### LIMITED WARRANTY

ECOSTADT TECHNOLOGIES LLC warrants that to the extent permitted by applicable law (a) this product conforms to chemical description on its label; (b) this product is reasonably fit for purposes stated on its label, subject to inherent risks referred to herein, when used in accordance with its directions; and (c) that directions, cautions and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. To the extent permitted by applicable law, there are no express warranties other than those set forth herein. ECOSTADT TECHNOLOGIES LLC neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. To the extent permitted by applicable law, ECOSTADT TECHNOLOGIES LLC expressly excludes and disclaims all implied warranties of merchantability, fitness for purpose (or) any other warranty of quality of performance.

To the extent permitted by applicable law, this warranty does not extend to, and user shall be solely responsible for, any loss or damage that results from the use of this product in any manner that is inconsistent with this label's directions, or cautions.

To the extent permitted by applicable law, user's exclusive remedy and ECOSTADT TECHNOLOGIES LLC's or seller's exclusive liability for any claim loss, damage, or injury resulting from use or handling of this product, whether (or) not based in contract, negligence, strict liability in tort, (or) otherwise, shall be limited, at ECOSTADT TECHNOLOGIES LLC's option, to replacement, (or) repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

#### [Optional marketing claims, may be used on any panel]

- Controls Listed Insects, Mites and Nematodes.
- Insecticide, Miticide & Nematicide
- Broad Spectrum Control.
- Can be used the day of harvest.
- Rose and flower spray.
- Garden vegetable and Ornamental spray.
- Tomato and vegetable spray.
- Foliar and systemic application.

[Note: Texts in [ ] are optional language]

### ECOWORKS EC (EPA Reg. #. 89152- 4)

Sub Label D: Residential/Home Owner Use

### **ECOWORKS EC**

[Additional Brand Names- ECOHALT EC, NEEM WORKS EC, ECOMAXIMA EC]
BOTANICAL INSECTICIDE, MITICIDE, NEMATICIDE & FUNGICIDE
READY TO MAKE PRODUCT (RTM)
PREMEASURED SINGLE DOSE WATER SOLUBLE SACHET

FOR HOME GARDEN AND RESIDENTIAL GREENHOUSE USE.
FOR THE CONTROL OF INSECTS, MITES, NEMATODES AND FUNGAL DISEASES.
PRODUCT CAN BE USED UP TO DAY OF HARVEST IN ALL FOOD AND NON-FOOD CROPS.





[\* Contains 2200 PPM Azadirachtin]

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### **FIRST AID**

#### If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

#### If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

#### If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- · Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call your poison control center at 1-800-222-1222. For information on product, usage, call 1-888-858-7378, Seven days a week, 6:30 am to 4.30 pm Pacific Time. During other times, call the poison control center 1-800-222-1222.

Medical: Hazard information Services (H.I. S1) 800-228-5635, Extension 169.

ECOSTADT TECHNOLOGIES LLC 1024 Iron Point Road, Suite 100-1318

Folsom, CA 95630 Phone: 916-357-6607 Net Contents: 5 sachets (Total Volume: 50ml, 3 1/3 Tbs), 10 sachets (Total Volume: 100ml, 6 2/3 Tbs), 20 sachets (Total Volume: 200ml, 13 1/3 Tbs), 40 sachets (Total Volume: 400ml,

26 2/3Tbs). Each sachet contains 10 ml (2/3 Tbs)

EPA Reg.# 89152-4

EPA Est. # Batch No.:

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before use.

#### **ENVIRONMENTAL HAZARDS**

For Terrestrial Uses Only. This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage, ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### PRODUCT INFORMATION

ECOWORKS EC is an emulsifiable concentrate containing 70% cold pressed Neem oil. ECOWORKS EC controls insects such as worms, thrips, aphids, white flies, leaf miners, beetles, mites and nematodes etc. ECOWORKS EC repels, disrupts feeding and growth of insect pests.

ECOWORKS EC is effective for the prevention and control of fungal diseases such as powdery mildew, downy mildew, rust, leaf spot, scab, blight, anthracnose, dollar spot, molds etc.

#### **APPLICATION INSTRUCTIONS**

#### READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USE

ECOWORKS EC is exempt from tolerances and may be applied as directed to any food or non-food crop up to and including day of harvest at a rate not exceeding 14 Pints (20 grams active ingredient) per acre per application.

#### **GENERAL PRECAUTIONS**

Do not apply ECOWORKS EC on just germinated, newly transplanted, wilted and drought stressed plants and plants with tender leaves.

As followed in oil based formulations, do not apply ECOWORKS EC during hot temperature conditions (>90 F). It is recommended to apply during early morning and late evening time to prevent potential leaf burns and plant injuries.

Exercise care when combining with oil and surfactant based products.

It is recommended not to apply to sensitive plant species such as poinsettias, some rose species, impatiens, hibiscus flowers, carnations, Comice pears etc.

Waxy bloom on certain ornamental plants may be reduced after an application.

Applications may remove 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

**MIXING AND STORAGE:** Shake well before mixing. Always use this product promptly after mixing with water. Use spray water at ambient conditions (20-35 or 68-95°F) for mixing and avoid using cold waters. In case spray water is cold, first premix the product with a few gallons of water at 20-35 or 68-95°F and then mix with cold water. Use spray mixture immediately or within few hours of mixing.

#### (1). ECOWORKS EC USES FOR HOME GARDEN PLANTS- FOLIAR APPLICATIONS

For foliar applications, apply ECOWORKS EC with hand operated or power sprayer equipment for low volume. Follow equipment manufacturer recommendations when making applications.

Open ECOWORKS EC packet containing single dose vials. Drop required number of vials in to hand sprayer as per table below, fill water, shake well and spray.

#### ECOWORKS EC FOLIAR APPLICATION RATES FOR HOME GARDEN PLANTS SUCH AS VEGETABLES, HERBS, SPICES, FRUITS, BERRIES AND ORNAMENTALS

|   | PESTS &  |                        | COWORI                  | APPLICATION            |   |
|---|--|------------------------|-------------------------|------------------------|---|
| CROP  |  | One<br>Gallon<br>water | Five<br>Gallon<br>water | Ten<br>Gallon<br>water | INTERVALS   |
| Leafy Vegetables including but not limited to: Broccoli, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew, | 1<br>sachet            | 5<br>sachets            | 10<br>sachets          | 7 - 14 days.  Spray early as soon as pest infection is noticed. |

|   | Downy mildew, rust  |             |              |               |   |
|---|---|-------------|--------------|---------------|---|
| Root Vegetables including but not limited to: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yam                               | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>rust | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Fruiting Vegetables including but not limited to: Eggplant, Pepper, Tomatillo, Tomato   | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites<br>Powdery<br>mildew,<br>Downy mildew,<br>rust  | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Cucurbit Vegetables including but not limited to: Cucumber, Gourd, Muskmelon, Pumpkin, Squash, Watermelon, Cantaloupe, Casaba, Gherkins, Melons, Zucchini | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites<br>Powdery<br>mildew,<br>Downy mildew,          | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Legume Vegetables including but not limited to: Bean, Chickpea, Lentil, Pea   | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites<br>Powdery<br>mildew,<br>Downy mildew,<br>rust  | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Bulb Vegetables including but not limited to:   | Leafrollers,<br>Loopers,<br>Armyworms,  | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.<br>Spray early as<br>soon as pest                  |

| Garlic, Onion, Shallot   | Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew,  |              |              |               | infection is noticed.   |
|--|---|--------------|--------------|---------------|---|
| Berries including but<br>not limited to:<br>Blackberry,<br>Blueberry,<br>Raspberry,<br>Strawberry,<br>Gooseberry                     | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Anthracnose, Botrytis   | 1<br>sachets | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Herbs and Spices including but not limited to: Chive, Dill, Fennel, Mustard, Sage, Basil, Coriander, Peppermint                      | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew, | 1<br>sachet  | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Nut trees including<br>but not limited to:<br>Almond, Walnut,<br>Pecan, Pistachio  | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew, | 1<br>sachet  | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Pome fruits including<br>but not limited to:<br>Apple, Pear (check<br>plant safety for<br>Comice Pear variety<br>before application) | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew, | 1<br>sachet  | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |

|  | Anthracnose,   |             |              |               |   |
|--|--|-------------|--------------|---------------|---|
|  | Botrytis   |             |              |               |   |
| Stone fruits including<br>but not limited to:<br>Apricot, Cherry,<br>Nectarine, Peach,<br>Plum | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Anthracnose, Botrytis                      | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Citrus fruits including<br>but not limited to:<br>Grapefruit, Lemon,<br>Orange                 | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Anthracnose, Botrytis                      | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7- 14 days. Spray early as soon as pest infection is noticed.   |
| Ornamental Plants including but not limited to: Chrysanthemum, Coleus, Roses, Poinsettia       | Leafrollers, Loopers, Armyworms, Whiteflies, Aphids, Thrips, Mites, Powdery mildew, Downy mildew, Blackspot, rust                            | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |
| Ornamental Trees & Shrubs  | Leafrollers,<br>Loopers,<br>Armyworms,<br>Whiteflies,<br>Aphids,<br>Thrips, Mites,<br>Powdery<br>mildew,<br>Downy mildew,<br>Blackspot, rust | 1<br>sachet | 5<br>sachets | 10<br>sachets | 7 - 14 days.  Spray early as soon as pest infection is noticed. |

# (2). ECOWORKS EC USES FOR HOME GARDEN PLANTS, INDOOR AND OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, INTERIORSCAPES, AND SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS- DRENCH APPLICATION

Use ECOWORKS EC as a soil drench for effective control of foliar and soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or thrips, fungal diseases, etc.

Open ECOWORKS EC packet containing single dose vials. Drop required number of vials in to bucket with water as per table below and stir bucket well.

#### ECOWORKS EC APPLICATION TABLE FOR HOME & GARDENS DRENCH APPLICATIONS FOR INDOOR AND OUTDOOR PLANTS RESIDENTIAL GREENHOUSES, INTERIORSCAPES, AND SOIL-LESS MEDIUM PLANTS GROWING SYSTEMS

| Gallons of | AM         | APPLICATION |            |               |             |  |  |
|------------|------------|-------------|------------|---------------|-------------|--|--|
| water      | 0.05%      | 0.1%        | 0.15%      | 0.2%          | INTERVAL    |  |  |
| 5 GALLONS  | 1 sachet   | 2 sachets   | 3 sachets  | 4 sachets     | 7 – 14 days |  |  |
| 10 GALLONS | 2 sachets  | 4 sachets   | 6 sachets  | 8 sachets     | 7 – 14 days |  |  |
| 20 GALLONS | 4 sachets  | 8 sachets   | 12 sachets | 16<br>sachets | 7 – 14 days |  |  |
| 30 GALLONS | 6 sachets  | 12 sachets  | 18 sachets | 24<br>sachets | 7 – 14 days |  |  |
| 40 GALLONS | 8 sachets  | 16 sachets  | 24 sachets | 32<br>sachets | 7 – 14 days |  |  |
| 50 GALLONS | 10 sachets | 20 sachets  | 30 sachets | 40<br>sachets | 7 – 14 days |  |  |

<sup>\*</sup>Drench soil in the pot with one (1) pint of finished product dilution per gallon of soil. Use lower concentration (0.05%) for mild infestations and higher concentration (0.2%) for severe infestation.

#### STORAGE AND DISPOSAL

#### Do Not Contaminate Water, Food, or Feed by Storage or Disposal.

Store in a cool dry place out of reach of children and pets and away from direct sunlight, windows, flames and other sources of heat or ignitions. Keep containers tightly closed and in original containers when not in use. Do not store product such that it is exposed to ultraviolet light (sunlight) or moisture.

Non-refillable container. Do not reuse or refill this container.

**If empty**: Place in trash or offer for recycling if available.

**If partly filled**: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### IMPORTANT: PLEASE READ BEFORE USE

#### LIMITED WARRANTY

ECOSTADT TECHNOLOGIES LLC warrants that to the extent permitted by applicable law (a) this product conforms to chemical description on its label; (b) this product is reasonably fit for purposes stated on its label, subject to inherent risks referred to herein, when used in accordance with its directions; and (c) that directions, cautions and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. To the extent permitted by applicable law, there are no express warranties other than those set forth herein. ECOSTADT TECHNOLOGIES LLC neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. To the extent permitted by applicable law, ECOSTADT TECHNOLOGIES LLC expressly excludes and disclaims all implied warranties of merchantability, fitness for purpose (or) any other warranty of quality of performance.

To the extent permitted by applicable law, this warranty does not extend to, and user shall be solely responsible for, any loss or damage that results from the use of this product in any manner that is inconsistent with this label's directions, or cautions.

To the extent permitted by applicable law, user's exclusive remedy and ECOSTADT TECHNOLOGIES LLC's or seller's exclusive liability for any claim loss, damage, or injury resulting from use or handling of this product, whether (or) not based in contract, negligence, strict liability in tort, (or) otherwise, shall be limited, at ECOSTADT TECHNOLOGIES LLC's option, to replacement, (or) repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

#### [Optional marketing claims, may be used on any panel]

- Controls Listed Insects, Mites and Nematodes.
- Insecticide, Miticide & Nematicide
- Broad Spectrum Control.
- Can be used the day of harvest.
- Rose and flower spray.
- Garden vegetable and Ornamental spray.
- Tomato and vegetable spray.
- Foliar and systemic application.